
# Drinking guidelines used in the context of early identification and brief interventions in Europe: overview of RARHA survey results

E. Scafato, C. Gandin, L. Galluzzo, S. Ghirini, S. Martire Istituto Superiore di Sanità, Italy









#### ISS Work Group and RARHA Italian National Team

#### RARHA NATIONAL TEAM MEMBERS and INSTITUTIONS

1	Bologna	Emanuela	ISTAT					
2	Burgio	Alessandra	ISTAT					
3	Ceccolini	Carla	Ministero della Salute					
4	Crialesi	Roberta	ISTAT					
5	Galluzzo	Lucia	Istituto Superiore di Sanità					
6	Gandin	Claudia	Istituto Superiore di Sanità					
7	Gargiulo	Lidia	ISTAT					
8	Ghirini	Silvia	Istituto Superiore di Sanità					
9	Ghiselli	Andrea	EX INRAN oggi CRA					
10	Loghi	Marzia	ISTAT					
11	Martire	Sonia	Istituto Superiore di Sanità					
12	Quattrociocchi	Luciana	ISTAT					
13	Sante	Orsini	ISTAT					
14	Scafato	Emanuele	Istituto Superiore di Sanità					
15	Spizzichino	Lorenzo	Ministero della salute					
16	Tamburini	Cristina	Ministero della salute					
17	Tinto	Alessandra	ISTAT					
18	Vichi	Monica	Istituto Superiore di Sanità					
19	Solipaca	Alessandro	ISTAT-Osservatorio sulle regioni					
20	Ricciardi	Walter	Osservatorio sulle regioni					
21	Carle	Flavia	Ministero della salute-SDO					
22	Migliore	Maria	Ministero della salute					









#### WP5: Outline of the work (tasks)



- 1. Overview of drinking guidelines given in MS and of their main features (ISS)
- 2. Overview of the use of drinking guidelines in the context of Early Identification and Brief Interventions (EIBI) on Hazardous/Harmful Alcohol Consumption (HHAC) in PHC and other settings, drawing in particular on projects ODHIN and BISTAIRS (ISS)
- 3. Overview of guidelines on drinking by young people (LWL)
- 4. Overview of science underpinnings drawing on recent work done for Australian and Canadian guidelines (THL)
- 5. Overview of "standard drink" definitions across the EU and of main approaches to increase awareness of such tools for monitoring alcohol consumption (HSE)
- 6. Mapping consumer views on risk/safety communication as an approach to reduce alcohol related harm by on-line surveys in 16 MS (EUROCARE)
- 7. From science to practice: Expert/policymaker meeting (at ISS) to discuss preliminary results and conclusions from the overviews and to help develop a policy Delphi survey (THL)
- 8. Second Expert/policymaker meeting to foster dialogue on good practice principles in the use of drinking guidelines as a public health measure drawing on all previous tasks
- 9. Coordination and production of synthesis report (THL)





#### **ISS Data collection**




- ✓ The questionnaire has been submitted by email (July 2014) to the country representatives of the Committee on National Alcohol Policy and Action CNAPA as experts with qualified experience and competence on the specific topic
- ✓ Participants were asked to:
  - check the validity of the information derived from a preliminary ISS review of sources of information on the use of low risk drinking guidelines in EIBI ("review of available sources")
  - provide the most updated and reliable information for their Country ("RARHA survey")
- ✓ Detailed instructions to fill in the Country report/Questionnaire, standardizing the data collection were provided
- ✓ The present version of this overview has been amended (May 2016) with all corrections suggested by respondents to the 2014 survey. Data on guidelines/recommendations issued after the data collection were not included.





### The RARHA survey questionnaire





The **sources** of available information are:











#### **PARTICIPATION**



- ✓ 31 European countries addressed

  (all RARHA associated and collaborating countries + 1 additional country\*).
- ✓ 30 out of 31 European countries replied (Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic\*, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Norway, The Netherlands, Poland, Portugal, Romania, Slovenia, Spain, Sweden, Switzerland, United Kingdom).
- ✓ Slovakia did not reply





#### **Drinking guidelines in EIBI context in EU countries**




Country	Source				Is there a formal governmental organization, or organization appointed/contracted by the government that is responsable for preparing clinical guidelines for managing HHAC?		Are there multidisciplinary guidelines for managing HHAC in your country that have been approved or endorsed by at least one health care professional body or scientific societies?			Guidelines or recommendations		
Country	NIHGO	BISTAIRS	WHO 2013	вавна	1= Yes	; 2=No ; 3=Incon	sistent	1= Yes ; 2=No ; 3=Inconsistent ; 4=Under preparation			for BI / Treatment	
AUSTRIA						2	2		2	2		
BELGIUM					2	3	2	1	1	1		
BULGARIA							2			2		
CROATIA					1		1	1		1		☑
CYPRUS					1		1	2		2		
CZECH REPUBLIC					2	3	2	1	1	1	D	☑
DENMARK						1	1		3	2	D	☑
ESTONIA					2		1	2		1	D	☑
FINLAND					1	1	1	1	1	1	D	☑
FRANCE							1			1		☑
GERMANY					2	1	2	1	1	1	V	☑
GREECE					2	1	1	2	2	1	D	☑
HUNGARY							1			1	D	☑
ICELAND					1		1	1		1		☑
IRELAND					1	1	1	1	1	1		☑
ITALY					1	1	1	1	1	1	D	☑
LATVIA					1		1	1		1		☑
LITHUANIA						2	2		1	1		☑
LUXEMBOURG							2			2	D	
MALTA					1		1	2		2		☑
NETHERLANDS (THE)					1	1	1	1	1	1		☑
NORWAY							3			2		
POLAND					2	3	1	2	3	1		☑
PORTUGAL					1	1	1	1	4	1	☑	☑
ROMANIA					2		2	2		2		
SLOVAKIA						2			2			
SLOVENIA					2		1	1		1		✓
SPAIN					1	1	1	1	1	1	V	☑
SWEDEN					1	3	1	1	1	1		☑
SWITZERLAND					2		2	1		1		
UNITED KINGDOM					1	1	1	1	1	1		☑






## 1) Formal governmental organization(or similar) responsible for clinical guidelines for managing HHAC






2) Multidisciplinary guidelines for managing HHAC in EU countries ...





#### 3) Guidelines or recommendations for BI / Treatment





#### **Conclusions**



## Drinking guidelines are used in the context of early identification and brief interventions in Europe:

- ✓ The number of organizations formally appointed to develop clinical guidelines for managing HHAC has increased over time (20/31)
- ✓ The large majority of investigated countries (22/31) has multidisciplinary guidelines for managing HHAC
- ✓ Guidelines or recommendations specific for BI/ treatment are available in 22/31 EU countries





## Drinking guidelines used in the context of early identification and brief interventions in Europe: overview of RARHA survey results

#### E. Scafato , C. Gandin, L. Galluzzo, S. Ghirini, S. Martire

Istituto Superiore di Sanità, Italy

E-mail: <a href="mailto:emanuele.scafato@iss.it">emanuele.scafato@iss.it</a> Website: <a href="mailto:www.iss.it">www.iss.it</a>





